

GEOMETRICITY AND POLYGONALITY IN FREE GROUPS

SANG-HYUN KIM

ABSTRACT. Gordon and Wilton recently proved that the double D of a free group F amalgamated along a cyclic subgroup C of F contains a surface group if a generator w of C satisfies a certain 3-manifold theoretic condition, called virtually geometricity. Wilton and the author defined the polygonality of w which also guarantees the existence of a surface group in D . In this paper, virtual geometricity is shown to imply polygonality up to descending to a finite-index subgroup of F . That the converse does not hold will follow from an example formerly considered by Manning.

[2000]Primary 20F36, 20F65 Whitehead graph, subgroup separability

1. INTRODUCTION

A *surface group* (*hyperbolic surface group*, respectively) means a group isomorphic to the fundamental group of a closed surface with non-positive (negative, respectively) Euler characteristic. Throughout this paper, we let F be a finitely generated non-abelian free group, and H be a handlebody so that $\pi_1(H) = F$. Unless specified otherwise, we fix $\mathcal{S} = \{a_1, \dots, a_n\}$ as a generating set for F . Each element $w \in F$ can be written as a *word* written in $\mathcal{S} \cup \mathcal{S}^{-1}$. That means, $w = v_1 v_2 \cdots v_l$ for some $v_i \in \mathcal{S} \cup \mathcal{S}^{-1}$. Each v_i is called a *letter* of w . w is *cyclically reduced* if $v_i \neq v_{i+1}^{-1}$, where the indices are taken modulo l . The length of a cyclically reduced word w is notated as $|w|$. The Cayley graph of F with respect to the generating set $\mathcal{S}$ is an infinite $2n$ -valent tree, denoted as $\text{Cay}_{\mathcal{S}}(F)$ or simply $\text{Cay}(F)$. There is a natural action of F on $\text{Cay}(F)$, so that $\text{Cay}(F)/F$ is a bouquet of n circles. Each circle in $\text{Cay}(F)/F$ inherits from $\text{Cay}(F)$ an orientation and a label by an element in $\mathcal{S}$.

Motivated by 3-manifold theory, Gromov conjectured every one-ended word-hyperbolic group contains a surface group [2]. Note that a surface subgroup of a word-hyperbolic group is actually a hyperbolic surface group. We say that $U \subseteq F$ is *diskbusting* if there does not exist a non-trivial free decomposition $F = G_1 * G_2$ such that each element of U is conjugate into one of G_i 's [5, 12, 10]. For convention, we will only consider finite subsets of F . As a special case, a word $w \in F$ is diskbusting if w does not belong to any proper free factor of F . $w \in F$ is *root-free* if w is not a proper power. A particularly simple case of the Gromov's conjecture is when a word-hyperbolic group is given as the *double* $D(w) = F *_{\langle w \rangle} F$. $D(w)$ is one-ended and word-hyperbolic if and only if w is diskbusting and root-free [3, 6]. Gordon and Wilton ignited attention on this case by providing several sufficient conditions for $D(w)$ to contain a surface group [6]. On one hand, they formulated a homological condition by crucially using a result of Calegari on surface subgroups of word-hyperbolic graphs of free groups with cyclic edge groups [4]. On the other hand, they considered a 3-manifold theoretic condition as follows. Realize $U \subseteq F = \pi_1(H)$ as an embedded 1-submanifold $A \subseteq H$. U (or equivalently, A) is said to be *virtually*

geometric if there exists a finite cover $p : H' \rightarrow H$ such that $p^{-1}(A)$ is freely homotopic to a 1-submanifold embedded in $\partial H'$. In particular, U (or equivalently, A) is *geometric* if A is freely homotopic to a 1-submanifold on ∂H . If $w \in F$ is virtually geometric and diskbusting, then $D(w)$ contains a surface group [6]. However, not all root-free diskbusting words are virtually geometric [8].

Let $w \in F = \pi_1(\text{Cay}(F)/F)$ be cyclically reduced. Following [11], a locally injective graph map is called an *immersion*. There exists an immersed loop $\gamma \subseteq \text{Cay}(F)/F$ such that $[\gamma] = w$; in this case, we say that γ *reads* w . Let $X(w)$ denote the 2-dimensional CW-complex obtained by taking two copies of $\text{Cay}(F)/F$ and gluing each boundary component of a cylinder along each copy of $\gamma_w \subseteq \text{Cay}(F)/F$. In the case when the free basis $\mathcal{S}$ needs to be explicitly notated, we write $X_{\mathcal{S}}(w) = X(w)$. It turns out that $X(w)$ is an Eilenberg–Mac Lane space for $D(w)$ [15, 6].

In [7], Wilton and the author defined a combinatorial condition for a cyclically reduced word w , called *polygonality*; see Definition 1. A key observation was, the polygonality of w is equivalent to the existence of a homeomorphically and π_1 -injectively embedded closed surface of non-positive Euler characteristic in some finite cover of $X(w)$. Hence, the polygonality of w will guarantee the existence of a surface group in $D(w)$.

We will use the following terminology that appears in [7]. A *polygonal disk* P is a 2-dimensional closed disk with a CW-structure such that $\partial P = P^{(1)}$ is a polygon. Let $E(\partial P)$ denote the set of the edges in ∂P . A *side-pairing* $\sim$ on a collection of polygonal disks $P_1, \dots, P_m$ is a partition of $\coprod_i E(\partial P_i)$ into unordered pairs, along with a choice of a homeomorphism between the edges in each pair; here, we require that such a homeomorphism does not identify two consecutive edges of any ∂P_i in a way that fixes a common vertex of the two edges. A side-pairing $\sim$ on polygonal disks $P_1, \dots, P_m$ determines an identification of the edges of $\coprod_i P_i$; hence, we have a closed surface $S = \coprod_i P_i / \sim$. We denote by $m(S)$ the number of polygonal disks (2-cells) which S is made of. If $\phi : \Gamma \rightarrow \text{Cay}(F)/F$ is a graph map, each $e \in E(\Gamma)$ carries an orientation and a label (by $\mathcal{S}$) induced from the orientation and the label of $\phi(e) \in E(\text{Cay}(F)/F)$. Conversely, for a graph Γ , a choice of an orientation and a label (by $\mathcal{S}$) for each $e \in E(\Gamma)$ determines a graph map $\phi : \Gamma \rightarrow \text{Cay}(F)/F$; if $e \in E(\Gamma)$ is labeled by a_i , we call e as an a_i -edge of Γ . That $\phi : \Gamma \rightarrow \text{Cay}(F)/F$ is an immersion is equivalent to that for each $a_i \in \mathcal{S}$ and each $v \in \Gamma^{(0)}$, there do not exist two incoming a_i -edges or two outgoing a_i -edges at v . We say that $U \subseteq F$ is *independent* if for any two distinct $w_1, w_2 \in U$ and for any $t_1, t_2 \in \mathbb{Z} \setminus \{0\}$, $w_1^{t_1}$ is not conjugate to $w_2^{t_2}$. The definition of polygonality generalized to an independent set of cyclically reduced words is as follows; see [7] for the case when $|U| = 1$.

Definition 1. Let $U \subseteq F$ be an independent set of cyclically reduced words.

- (1) Suppose there exists a side-pairing $\sim$ on polygonal disks $P_1, \dots, P_m$ and an immersion $\phi : S^{(1)} \rightarrow \text{Cay}(F)/F$ where $S = \coprod_i P_i / \sim$, such that the composition $\partial P_i \rightarrow S^{(1)} \rightarrow \text{Cay}(F)/F$ reads a non-trivial power of some word in U . Then the closed surface S is called a *U -polygonal surface*.
- (2) U is *polygonal* if either U contains a proper power or there exists a U -polygonal surface S such that $\chi(S) < m(S)$.

Remark 2. Let $U \subseteq F$ be a set of cyclically reduced words. The condition $\chi(S) < m(S)$ is necessary for the polygonality to be a non-trivial property, for one can always construct a U -polygonal surface S with $\chi(S) = m(S)$ as follows. Choose

any non-trivial word $w \in U$ and take two polygonal disks P and P' each of whose boundary components reads w . For $e \in E(\partial P)$ and $e' \in E(\partial P')$, define $e \sim e'$ if and only if e and e' read the same letter of w . Then for $S = P \amalg P' / \sim$, $\chi(S) = m(S) = 2$. One can also construct a U -polygonal surface $S = P / \sim$ with $\chi(S) = m(S) = 1$ by taking a polygonal disk P whose boundary reads w^2 and by defining a side-pairing $\sim$ to identify the edges of ∂P with respect to the π -rotation.

The polygonality of $U \subseteq F$ depends on the choice of a free basis $\mathcal{S}$ for F in which U is written. For $g, h \in F$, we let g^h denote $h^{-1}gh$. Choose a finite-index subgroup F' of F and a free basis $\mathcal{S}'$ for F' . For $w' \in F'$, let us denote the F' -conjugacy class of w' by $[w']$. Fix a word $w \in F$. For $g \in F$, let n_g be the smallest positive integer such that $(w^{n_g})^g \in F'$. Following [6, 8], define

$$\underline{w}_{F'} = \{[(w^{n_g})^g] \mid gF' \in F/F'\}.$$

Note that $|\underline{w}_{F'}| \leq |F/F'|$. We define a *transversal* $\hat{w}_{F'}$ for $\underline{w}_{F'}$ to be a subset of F' obtained by choosing exactly one element from each conjugacy class in $\underline{w}_{F'}$; here, we will regard $\hat{w}_{F'}$ as a set of cyclically reduced words written in $\mathcal{S}'$, by taking cyclic conjugations if necessary. Suppose for some g and h in F , $(w^{n_g})^g$ and $(w^{n_h})^h$ have non-trivial powers which are conjugate to each other in F' . Since F does not have any non-trivial Baumslag–Solitar relation, we can write $(w^M)^g = (w^M)^{hf'}$ for some $M > 0$ and $f' \in F'$. We have $hf'g^{-1} \in C(w^M) = C(w)$ and so, $w^{hf'} = w^g$. In particular, $n_g = n_h$ and $[(w^{n_g})^g] = [(w^{n_h})^h]$. This shows that $\hat{w}_{F'}$ is independent as a set of words in F' .

Definition 3. A word $w \in F$ is *virtually polygonal* if a transversal $\hat{w}_{F'}$ for $\underline{w}_{F'}$ is polygonal as an independent set of cyclically reduced words written in $\mathcal{S}'$, for some finite-index subgroup F' of F and for some free basis $\mathcal{S}'$ of F' .

After formulating polygonality in terms of Whitehead graphs, we will prove a relation between virtual geometricity and virtual polygonality.

Theorem (Theorem 25). *A diskbusting and virtually geometric word in F is virtually polygonal.*

While Theorem 25 is interesting in its own, it also has a corollary related to the Gromov’s conjecture on $D(w)$.

Theorem (Theorem 26). *If a root-free word $w \in F$ is virtually polygonal, then $D(w)$ contains a hyperbolic surface group.*

Corollary (Corollary 27; first proved by Gordon–Wilton [6]). *If $w \in F$ is root-free, virtually geometric and diskbusting, then $D(w)$ contains a hyperbolic surface group.*

In [8], Manning proved that the word $w_1 = bbaaccabc \in F_3 = \langle a, b, c \rangle$ is not virtually geometric. The same argument shows that $w_2 = aabbacbccadbdcd \in F_4 = \langle a, b, c, d \rangle$ is not virtually geometric. We will prove that w_1 and w_2 are both polygonal (Proposition 32). As a consequence, $D(w_1)$ and $D(w_2)$ contain surface groups.

Theorem (Theorem 28). *There exist polygonal words which are not virtually geometric.*

2. PRELIMINARY

We recall basic facts on polygonality and Whitehead graphs. The material in Section 2.1 is largely drawn from [7]. For readers' convenience, we include a complete proof of Theorem 6 which is similar to the argument in [7, Corollary 2.11].

2.1. Polygonality. We say that $U, U' \subseteq F$ are *equivalent* if $U' = \phi(U)$ for some $\phi \in \text{Aut}(F)$. The definition of polygonality of $U \subseteq F$ involves the representation of U as a set of words written in $\mathcal{S} = \{a_1, \dots, a_n\}$ (Definition 1). In particular, $U \subseteq F$ is equivalent to a polygonal set of words if and only if $U \subseteq F$ is polygonal with respect to some choice of a free basis for F . We list some of the known polygonal words as follows.

Example 4. (1) Consider a cyclically reduced word $w \in F = \langle a_1, \dots, a_n \rangle$ such that for each i , exactly two letters of w belong to $\{a_i, a_i^{-1}\}$. In particular, $|w| = 2n$. We claim that w is polygonal. We may assume w is root-free, as any proper powers are polygonal by definition. Let P be a polygonal disk with each edge oriented and labeled by $\mathcal{S}$ such that ∂P reads w . Let $\sim$ be the side-pairing which identifies the two a_i -edges of ∂P for each i . Put $S = P/\sim$. Since $S^{(1)}$ has exactly one a_i -edge for each i , $S^{(1)}$ immerses into $\text{Cay}(F)/F$. In order to prove w is polygonal, we have only to show that $\chi(S) < m(S) = 1$. For each $v \in S^{(0)}$, let d_v denote the valence (of the 1-complex $S^{(1)}$) at v . As w is cyclically reduced, $d_v \geq 2$ for each v . Moreover, $\chi(S) = |S^{(0)}| - |S^{(1)}| + 1 = \sum_{v \in S^{(0)}} 1 - \sum_{v \in S^{(0)}} d_v/2 + 1$. Hence, $\chi(S) < 1$ fails only when $d_v = 2$ for each $v \in S^{(0)}$. If $d_v = 2$ for each v , then one can see that $S \approx \mathbb{R}P^2$ and $\sim$ identifies the edges of ∂P by the π -rotation; this means, $\sim$ identifies v_i to $v_{(|w|/2)+i}$ for each i . It follows that $w = u^2$ for some $u \in F$, and we have a contradiction.

(2) Let F_2 denote the free group generated by a and b . If $|p_i|, |q_i| > 1$ for each i , then the words $\prod_i a^{p_i} b^{q_i}, \prod_i a^{p_i} (a^b)^{q_i} \in F_2$ are polygonal [7].

(3) The word $w = \prod_i a^{p_i} (a^b)^{q_i}$ is called a *positive height-1 word* of F_2 if $p_i, q_i > 0$. Put $p = \sum_i p_i, q = \sum_i q_i, p' = \sum_{p_i=1} 1$ and $q' = \sum_{q_i=1} 1$. If $pp' \leq q^2$ and $qq' \leq p^2$, then w is polygonal [7]. With a suitable notion of probability on F_2 , this implies a positive height-1 word is “almost surely” polygonal.

Let $p : (X', x') \rightarrow (X, x)$ be a finite covering map for some based spaces (X', x') and (X, x) . For a based loop $\gamma : (S^1, v) \rightarrow (X, x)$, let $q : (S^1, v') \rightarrow (S^1, v)$ be the smallest covering such that $\gamma \circ q$ lifts to a based loop $\tilde{\gamma} : (S^1, v') \rightarrow (X', x')$ as shown in the commutative diagram below. Following [15], we say that $\tilde{\gamma}$ is the *elevation of γ at x' with respect to p* .

$$\begin{array}{ccc} (S^1, v') & \xrightarrow{\tilde{\gamma}} & (X', x') \\ \downarrow q & & \downarrow p \\ (S^1, v) & \xrightarrow{\gamma} & (X, x) \end{array}$$

Let $U = \{w_1, \dots, w_r\} \subseteq F$ be an independent set of root-free, cyclically reduced words, and γ_i be the (based) loop in $\text{Cay}(F)/F$ reading w_i . Take two copies Γ_1, Γ_2 of $\text{Cay}(F)/F$ and glue each boundary component of a cylinder $C_i \approx S^1 \times [-1, 1]$ along each copy of γ_i in Γ_1 and in Γ_2 , for $i = 1, 2, \dots, r$. We let $X(U)$ denote the 2-dimensional CW-complex thus obtained. Define $D(U) = \pi_1(X(U))$. Take

$F^{(1)}, F^{(2)} \cong F$, and let $w_i^{(j)}$ denote the image of w_i in $F^{(j)}$ for $j = 1, 2$. Note that

$$D(U) \cong \langle F^{(1)}, F^{(2)}, t_2, t_3, \dots, t_r \mid w_1^{(1)} = w_1^{(2)}, (w_i^{(1)})^{t_i} = (w_i^{(2)}) \text{ for } i = 2, \dots, r \rangle.$$

For a finite-index subgroup $F' \leq F$, $\text{Cay}(F)/F'$ is the finite covering space of $\text{Cay}(F)/F$ corresponding to F' . For each choice of the basepoint $\tilde{x}$ of $\text{Cay}(F)/F'$, there is an elevation of γ_i at $\tilde{x}$ with respect to the covering $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$. We identify two elevations $\tilde{\gamma}, \tilde{\gamma}'$ (at two distinct basepoints $\tilde{x}, \tilde{x}' \in \text{Cay}(F)/F'$) of the based loop $\gamma_i: S^1 \rightarrow \text{Cay}(F)/F$ if $\tilde{\gamma}$ and $\tilde{\gamma}'$ are the same as loops without basepoints (*i.e.*, when the basepoints are forgotten); this means, the lifting of some power of γ_i at $\tilde{x}$ to $\text{Cay}(F)/F'$ terminates at $\tilde{x}'$. See [14, Lemma 2.7] and [7, Lemma 2.4] for algebraic description of this identification. We let $\{\tilde{\gamma}_1, \tilde{\gamma}_2, \dots, \tilde{\gamma}_s\}$ denote the set of all the elevations of $\gamma_1, \gamma_2, \dots, \gamma_r$ at the vertices of $\text{Cay}(F)/F'$, after this identification. Take two copies Γ'_1, Γ'_2 of $\text{Cay}(F)/F'$ and glue each boundary component of a cylinder C'_j along each copy of $\tilde{\gamma}_j$ in Γ'_1 and in Γ'_2 , for $j = 1, 2, \dots, s$; in this way, one obtains a finite cover $Y(U, F')$ of $X(U)$. The image of each cylinder in $X(U)$ or in $Y(U, F')$ will still be called a *cylinder*. Lemma 5 is a special case of [7, Lemma 2.2]. We omit the proof here.

Lemma 5 ([7]). *Suppose $U \subseteq F$ is an independent set of root-free, cyclically reduced words and $[F : F'] < \infty$. Let S be a closed connected surface homeomorphically embedded in $Y(U, F')$. Then S is the union of some cylinders, $\chi(S) \leq 0$ and $\pi_1(S)$ embeds into $D(U)$.* $\square$

The following generalizes [7, Corollary 2.11].

Theorem 6. *Let $U \subseteq F$ be an independent set of root-free, cyclically reduced words. If U is polygonal, then $D(U)$ contains a hyperbolic surface group.*

Proof. Write $U = \{w_1, \dots, w_r\}$, and let $\gamma_j \subseteq \text{Cay}(F)/F$ realize w_j . Let $\sim$ be a side-pairing on polygonal disks $P_1, \dots, P_m$ such that $S = \coprod_i P_i / \sim$ is a closed U -polygonal surface satisfying $\chi(S) < m$. This implies that for some immersion $\phi: S^{(1)} \rightarrow \text{Cay}(F)/F$, each composition $\partial P_i \rightarrow S^{(1)} \rightarrow \text{Cay}(F)/F$ reads a non-trivial power of an element in U . Since F is subgroup separable, one can lift the immersion ϕ to an embedding $\phi': S^{(1)} \rightarrow \text{Cay}(F)/F'$ for some $[F : F'] < \infty$ [9]. Choose an open disk B_i in the interior of each P_i and let S' be the double of $S \setminus \cup_i B_i$. Since ϕ' maps each ∂P_i to an elevation of some γ_j , the definition of $Y(U, F')$ implies that S' is homeomorphic to the union S'' of some cylinders in $Y(U, F')$. By Lemma 5, $\pi_1(S')$ embeds into $D(U)$. Note that $\chi(S') = 2(\chi(S) - m) < 0$. $\square$

Remark 7. Let $U \subseteq F$ be as in the hypothesis of Theorem 6. By an elementary argument on graphs of spaces, a finite cover of $X(U)$ contains a homeomorphically embedded closed surface if and only if so does $Y(U, F')$ for some $[F : F'] < \infty$. Then Theorem 6 can actually be strengthened as follows: U is polygonal if and only if a finite cover of $X(U)$ contains a homeomorphically embedded closed hyperbolic surface; see [7].

2.2. Whitehead graph. A *graph* means a 1-dimensional CW-complex. For a graph Γ , let $V(\Gamma)$ and $E(\Gamma)$ denote the vertex set and the edge set, respectively. For a cyclically reduced word $w = v_1 v_2 \dots v_l \in F$ where $v_i \in \mathcal{S} \cup \mathcal{S}^{-1}$, we let $W(w)$ denote the *Whitehead graph* of w . This means, $W(w)$ is a graph with the vertex set $\mathcal{S} \cup \mathcal{S}^{-1}$ and the edge set $\{e_1, e_2, \dots, e_l\}$ such that e_i joins v_i and v_{i+1}^{-1} , where the indices are taken modulo l . In the special case when $w = a_i$ or $w = a_i^{-1}$, $W(w)$

consists of a single edge joining a_i and a_i^{-1} . We define the *connecting map* σ_w associated with $W(w)$ as the map $\sigma_w: \{(e, v) | e \in E(W(w)), v \in \partial e\} \rightarrow E(W(w))$ such that $\sigma_w(e_i, v_i) = e_{i-1}$ and $\sigma_w(e_i, v_{i+1}^{-1}) = e_{i+1}$. In particular, if $\sigma(e, v) = e'$ then $v \in \partial e$, $v^{-1} \in \partial e'$ and moreover, the length-2 subwords of w corresponding to e and e' are consecutive and share the letter v or v^{-1} ; see Example 8.

Consider a non-zero integral polynomial $f(x_1, \dots, x_r) = \sum_{1 \leq i \leq r, 1 \leq j \leq s} c_{ij} x_i^j$ where $s > 0$ and $c_{ij} \geq 0$. For $U = \{w_1, \dots, w_r\} \subseteq F$ consisting of cyclically reduced words, we define

$$W(f(U)) = W(f(w_1, \dots, w_r)) = \cup_{i=1}^r \cup_{j=1}^s \cup_{k=1}^{c_{ij}} W(w_i^j)$$

where the union is taken so that each term $W(w_i^j)$ has the common vertex set $\mathcal{S} \cup \mathcal{S}^{-1}$, and any two terms do not have a common edge. For instance, $W(w + w)$ is obtained by doubling all the edges of $W(w)$, so that $V(W(w + w)) = \mathcal{S} \cup \mathcal{S}^{-1}$ and $|E(W(w + w))| = 2|E(W(w))| = 2|w|$. Note that $W(f(U))$ is an abusive notation in that U is considered as an *ordered tuple* rather than just a set. We have $|E(W(f(U)))| = \sum_{i,j} c_{ij} |E(W(w_i^j))| = \sum_{i,j} c_{ij} j |w_i|$. The unique extension on $W(f(U))$ of the connecting map on each $W(w_i^j)$ is called the *connecting map associated with* $W(f(U))$.

Example 8. . Consider the word $w = ab^{-2} \in F_2 = \langle a, b \rangle$. Write $w = v_1 v_2 v_3$ where $v_1 = a$ and $v_2 = v_3 = b^{-1}$. $W(w)$ is shown in Fig. 1 (a); here, the edges e_1, e_2 and e_3 correspond to the (cyclic) subwords $v_1 v_2 = ab^{-1}, v_2 v_3 = b^{-2}$ and $v_3 v_1 = b^{-1}a$, respectively. Some of the values of the associated connecting map are given as $\sigma_w(e_1, a) = \sigma_w(e_1, v_1) = e_3$, $\sigma_w(e_3, a^{-1}) = \sigma_w(e_3, v_1^{-1}) = e_1$ and $\sigma_w(e_3, b^{-1}) = \sigma_w(e_3, v_3) = e_2$.

Figure 1 (b) shows $W(w + w) = W(2w)$. e_i and e'_i denote the edges corresponding to the length-2 cyclic subword $v_i v_{i+1}$ of w . The connecting map is given as $\sigma_{2w}(e_1, a) = e_3$, $\sigma_{2w}(e_3, a^{-1}) = e_1$, $\sigma_{2w}(e_3, b^{-1}) = e_2$, $\sigma_{2w}(e'_1, a) = e'_3$ and so forth.

$W(w^2)$ is shown in Fig. 1 (c). f_i denotes the edge corresponding to the i -th length-2 cyclic subword of $w^2 = ab^{-2}ab^{-2}$. The connecting map can be computed as $\sigma_{w^2}(f_1, a) = f_6$, $\sigma_{w^2}(f_3, a^{-1}) = f_4$, $\sigma_{w^2}(f_3, b^{-1}) = f_2$, $\sigma_{w^2}(f_4, a) = f_3$ and so forth. Note that $W(2w)$ and $W(w^2)$ are the same as graphs, while σ_{2w} and σ_{w^2} are distinct.




FIGURE 1. $w = ab^{-2}$.

Remark 9. (1) Let σ be the connecting map associated with a Whitehead graph Γ . For $e \in E(\Gamma)$ and $v \in \partial e$, v^{-1} is an endpoint of $\sigma(e, v)$ and $\sigma(\sigma(e, v), v^{-1}) = e$.

(2) Let $U = \{w_1, \dots, w_r\}$ and $f(x_1, \dots, x_r) = \sum_{i,j \geq 1} c_{ij} x_i^j$. For each i and j , glue c_{ij} copies of a disk to $\text{Cay}(F)/F$ along the loop reading w_i^j . Let $Z(f(U))$ be the 2-dimensional CW-complex thus obtained. Then $W(f(U))$ is precisely the link of the (unique) vertex v in $Z(f(U))$. The vertex a_i (a_i^{-1} , respectively) of $W(f(U))$ corresponds to the incoming (outgoing, respectively) portion of the a_i -edge in $Z(f(U))^{(1)}$ at v .

A collection $\mathcal{D} = \{D_1, \dots, D_n\}$ of disjoint, properly embedded disks in H is called a *disk structure on H* if $H \setminus \cup_i D_i$ is a 3-cell [10]. We will always equip each disk D_i with a transverse orientation. Let A be an embedded 1-submanifold of H . Assume that A intersects $\mathcal{D}$ *transversely and minimally*. This means, $A \cap (\cup_i D_i)$ consists of finitely many points and $|A \cap (\cup_i D_i)| \leq |A' \cap (\cup_i D_i)|$ for any 1-submanifold A' freely homotopic to A . Given a disk structure $\mathcal{D}$ on H , we denote by $\mathcal{D} \times [-1, 1] = \cup_i (D_i \times [-1, 1])$ a closed regular neighborhood of $\mathcal{D}$ in H so that D_i is identified with $D_i \times 0$ and the transverse orientation of D_i is from $D_i \times -1$ to $D_i \times 1$. By transversality, we may assume that $A \cap (D_i \times [-1, 1])$ has the product structure $(A \cap D_i) \times [-1, 1]$ for each i . We define the *Whitehead graph $W(A, \mathcal{D})$ of A with respect to $\mathcal{D}$* as the graph such that the vertex set is $\mathcal{S} \cup \mathcal{S}^{-1}$ and the edge set consists of arcs in $A \setminus (\mathcal{D} \times (-1, 1))$, where $A \cap (D_i \times \pm 1)$ are identified with the vertices $a_i^{\pm 1}$ in $W(A, \mathcal{D})$. In other words, $W(A, \mathcal{D})$ is the graph obtained from $(A \cup (\mathcal{D} \times [-1, 1])) \setminus \mathcal{D} \times (-1, 1)$ by collapsing each $D_i \times \{\pm 1\}$ onto the vertices $a_i^{\pm 1}$. Suppose each loop γ_j in $A = \gamma_1 \cup \dots \cup \gamma_r$ is equipped with an orientation. Follow each loop γ_j , and whenever γ_j intersects D_i , record a_i or a_i^{-1} according to whether the orientation of γ_j coincides with the transverse orientation of D_i or not. Let w_j be the word thus obtained. In this case, we say that $U = \{w_1, \dots, w_r\}$ is *realized by A with respect to $\mathcal{D}$* . As we are assuming that A intersects $\mathcal{D}$ minimally, U consists of cyclically reduced words. Depending on the choice of the basepoint of γ_j , w_j is determined up to cyclic conjugation. Observe that $W(A, \mathcal{D}) = W(U)$, and the notion of the connecting map is also defined for $W(A, \mathcal{D})$ by that of $W(U)$. Moreover, for any other choice of disk structure $\mathcal{D}'$ on H which A intersects transversely and minimally, $W(A, \mathcal{D}') = W(\phi(U))$ for some $\phi \in \text{Aut}(F)$.

For a simple loop γ embedded in H and $c, j > 0$, we let $c\gamma^j$ denote the 1-submanifold consisting of c components, each of which is freely homotopic to γ^j . Consider a non-zero integral polynomial $f(x_1, \dots, x_r) = \sum_{i,j \geq 1} c_{ij} x_i^j$ where $c_{ij} \geq 0$. Let A be a 1-submanifold of H written as the union of disjoint loops $A = \gamma_1 \cup \dots \cup \gamma_r$. Then $f(A) = f(\gamma_1, \dots, \gamma_r)$ denotes a 1-submanifold in H freely homotopic to $\coprod_{i,j \geq 1} c_{ij} \gamma_i^j$. $f(A)$ is again an abusive notation in that A is considered as an ordered tuple of loops, rather than just a set of loops. If a 1-submanifold $A \subseteq H$ realizes $U \subseteq F$ with respect to a disk structure $\mathcal{D}$, then $W(f(U)) = W(f(A), \mathcal{D})$.

Remark 10. When considering the Whitehead graph of $U \subseteq F$ or a 1-submanifold $A \subseteq H$, we always require that U consists of cyclically reduced words, and A intersects $\mathcal{D}$ transversely and minimally. If necessary, we achieve this by replacing words in U by some cyclic conjugations or by freely homotoping loops in A . Note that certain additional conditions on A , such as $A \subseteq \partial H$, can possibly be lost by freely homotoping A .

A properly embedded disk D in H is *essential* if either $H \setminus D$ is connected or neither of the components of $H \setminus D$ is a 3-cell. Let A be a 1-submanifold of H . Suppose that for any 1-submanifold A' freely homotopic to A , A' intersects any essential disk $D \subseteq H$. Then A is said to be *diskbusting*. Note that a set U of cyclically reduced words in F is diskbusting if and only if a 1-submanifold $A \subseteq H$ realizing U is diskbusting.

A vertex v in a graph Γ is a *cut vertex* if $\Gamma \setminus \{v\}$ is not connected. We say that a Whitehead graph $W(A, \mathcal{D})$ is *minimal*^a if $|E(W(A, \mathcal{D}))| \leq |E(W(A', \mathcal{D}'))|$ for any 1-submanifold A' freely homotopic to A and any disk structure $\mathcal{D}'$ which A' intersects transversely and minimally. $U \subseteq F$ is *minimal* if U is not equivalent to any $U' \subseteq F$ where the sum of the lengths of the words in U' is smaller than that of U . Using a well-known finite-time algorithm to determine whether $U \subseteq F$ is diskbusting or not, one gets the following:

Theorem 11 ([13, 12, 10]). (1) *If $U \subseteq F$ is minimal, independent and diskbusting, then $W(U)$ is connected and does not have a cut vertex.*
 (2) *Suppose $A \subseteq H$ is a diskbusting 1-submanifold which intersects a disk structure $\mathcal{D}$ on H transversely and minimally, such that $W(A, \mathcal{D})$ is minimal. Then $W(A, \mathcal{D})$ is connected and does not have a cut vertex.* $\square$

3. SIMPLE SURGERY

Zieschang proved a key fact on a minimal Whitehead graph of a geometric 1-submanifold in H .

Theorem 12 ([16]; see also [1]). *Suppose A is a geometric 1-submanifold of H which intersects a disk structure $\mathcal{D}$ transversely and minimally. If $W(A, \mathcal{D})$ is minimal, then A is freely homotopic to some 1-submanifold $A' \subseteq \partial H$ such that A' intersects $\mathcal{D}$ transversely and minimally. In particular, $W(A', \mathcal{D}) = W(A, \mathcal{D})$.* $\square$

A *pairing* $\sim$ on a finite set X will mean an equivalence relation on X such that each equivalence class consists of precisely two elements. From now on, we will use the notation $I_1 = (0, 1]$ and $I_{-1} = [-1, 0)$.

Setting 13 (for Definition 14). Let $A \subseteq H$ be a 1-submanifold and $\mathcal{D} = \{D_1, \dots, D_n\}$ be a disk structure on H such that A intersects $\mathcal{D}$ transversely and minimally. Recall our convention that $\mathcal{D} \times [-1, 1] = \cup_i (D_i \times [-1, 1])$ denotes a closed regular neighborhood of $\mathcal{D}$ equipped with a product structure so that D_i is identified with $D_i \times 0$. In particular, if $v \in D_i$, then $v \times 1 \in D_i \times 1 \subseteq H$ and $v \times -1 \in D_i \times -1 \subseteq H$. We will also assume that A is compatible with the product structure in the sense that $A \cap (D_i \times [-1, 1]) = (A \cap D_i) \times [-1, 1]$ for each i . For each i , suppose $|A \cap D_i|$ is even and $\sim_i$ is a pairing on $A \cap D_i$. For each pair $v \sim_i v'$, let $\alpha_{vv'}^+ \subseteq D_i \times I_1$ be an arc joining $v \times 1$ and $v' \times 1$, and $\alpha_{vv'}^- \subseteq D_i \times I_{-1}$ be an arc joining $v \times -1$ and $v' \times -1$; moreover, we assume that all the arcs in $\{\alpha_{vv'}^\pm \mid v \sim_i v' \text{ for some } i\}$ are disjoint (see Fig. 2 and 3). Let $A' \subseteq H \setminus \mathcal{D}$ be the 1-submanifold obtained from A if one replaces $A \cap (D_i \times [-1, 1])$ by the arcs $\cup_{v \sim_i v'} \alpha_{vv'}^\pm$ for all i .

Definition 14. In Setting 13, assume further the following:

- (i) for each $1 \leq i \leq n$ and $\epsilon = \pm 1$, the intersection between each component of A' and $D_i \times I_\epsilon$ is either connected or empty;

^aMinimality of $W(A, \mathcal{D})$ should not be confused with the hypothesis that A intersects $\mathcal{D}$ transversely and minimally.

(ii) for some component γ of A' , $\gamma \cap (\mathcal{D} \times [-1, 1])$ has more than two components. Then we say that $(\mathcal{D}, \{\sim_i\})$ *determines a simple surgery on A* . When $\sim_i$ needs not be explicitly notated, we simply say A *admits a simple surgery (with respect to $\mathcal{D}$)*. Also, A' is said to be *obtained by the simple surgery $(\mathcal{D}, \{\sim_i\})$ on A* .

Example 15. Let $A \subseteq H$ be the 1-submanifold denoted as the bold curve in Fig. 2 (a). Write $D_1 \cap A = \{v, v'\}$ and $D_2 \cap A = \{w, w'\}$. Define $v \sim_1 v'$ and $w \sim_2 w'$, and consider $\alpha_{vv'}^\pm$ and $\alpha_{ww'}^\pm$ as in Setting 13. Put $\mathcal{D} = \{D_1, D_2\}$. Figure 2 (b) shows $A' \subseteq H$ which is obtained by substituting $\alpha_{vv'}^\pm$ and $\alpha_{ww'}^\pm$ for $A \cap (\mathcal{D} \times [-1, 1])$. Note that $A' \cap (\mathcal{D} \times [-1, 1])$ has four components $\{\alpha_{vv'}^+, \alpha_{vv'}^-, \alpha_{ww'}^+, \alpha_{ww'}^-\}$. It follows that A admits the simple surgery $(\mathcal{D}, \{\sim_1, \sim_2\})$.





FIGURE 2. A' is obtained by a simple surgery on A .

Example 16. The bold curves in Fig. 3 (a) denotes $A \subseteq H$. Put $D_1 \cap A = \{u, u', v, v'\}$ and $D_2 \cap A = \{w, w'\}$ as shown in the figure. Set $\mathcal{D} = \{D_1, D_2\}$.


Define $u \sim_1 u', v \sim_1 v'$ and $w \sim_2 w'$. Replace suitable segments on A by $\alpha_{uu'}^\pm, \alpha_{vv'}^\pm$ and $\alpha_{ww'}^\pm$ as in Setting 13, to obtain $A' = \gamma_1 \amalg \gamma_2 \amalg \gamma_3$; see Fig. 3 (b). Note that condition (ii) of Definition 14 is violated, since $\gamma_i \cap (\mathcal{D} \times [-1, 1])$ has exactly two components for each $i = 1, 2, 3$.

Consider a different pairing $u \sim_1 v', u' \sim_1 v$ and $w \sim_2 w'$. Following Setting 13, replace segments in $A \cap (\mathcal{D} \times [-1, 1])$ accordingly to obtain $A'' \approx S^1$ described in Fig. 3 (c). Condition (i) of Definition 14 then fails, since $A'' \cap (D_1 \times I_1)$ and


$A'' \cap (D_1 \times I_{-1})$ are both disconnected. In this way, one can see that A does not admit any simple surgery with respect to $\mathcal{D}$.



(a) $A \approx S^1 \amalg S^1 \amalg S^1 \amalg S^1$.



(b) $A' \approx S^1 \amalg S^1 \amalg S^1$.



(c) $A'' \approx S^1$.

FIGURE 3. A does not admit a simple surgery with respect to $\mathcal{D} = \{D_1, D_2\}$.

Whether $A \subseteq H$ admits a simple surgery or not can be detected by looking at a corresponding Whitehead graph.

Proposition 17. *Let A be a 1-submanifold of H which intersects a disk structure $\mathcal{D}$ transversely and minimally. Let σ denote the connecting map associated with*

$\Gamma = W(A, \mathcal{D})$. Then A admits a simple surgery with respect to $\mathcal{D}$ if and only if $\Gamma = \cup_h C_h$ for some C_h 's satisfying the following:

- (i) each C_h is a simple cycle,
- (ii) C_h and $C_{h'}$ do not have a common edge whenever $h \neq h'$,
- (iii) if e and e' are edges of some C_h intersecting at a vertex v , then $\sigma(e, v)$ and $\sigma(e', v)$ are edges of some $C_{h'}$,
- (iv) at least one C_h is not a bigon.

In condition (iii), h and h' can possibly be the same.

Proof. Write $\mathcal{D} = \{D_1, \dots, D_n\}$. We let $\Phi: (A \cup (\mathcal{D} \times [-1, 1])) \setminus (\mathcal{D} \times (-1, 1)) \rightarrow \Gamma$ denote the map collapsing $D_i \times \pm 1$ onto the vertices $a_i^{\pm 1}$.

($\Rightarrow$) Let A' be obtained from A by the simple surgery $(\mathcal{D}, \{\sim_i\})$. Φ maps each loop in A' onto a simple cycle in Γ , by the definition of a simple surgery. Hence, we have the condition (i). Let $\{C_1, C_2, \dots, C_t\}$ denote the set of simple cycles in Γ thus obtained. (ii) follows from the fact that A' consists of disjoint loops. For (iii), let $e, e' \in E(C_h)$ intersect with, say, $a_i \in V(\Gamma)$. Then there exist two arcs α, α' in $A \setminus (\mathcal{D} \times (-1, 1))$ and $v, v' \in A \cap D_i$ such that $v \sim_i v'$, $v \times 1 \in \alpha$, $v' \times 1 \in \alpha'$ and, e and e' are the images of α and α' by Φ . Let f and f' be the edges in Γ which are the images of the arcs in $A \setminus (\mathcal{D} \times (-1, 1))$ intersecting the disk $D_i \times -1$ at $v \times -1$ and at $v' \times -1$, respectively. By the definition of σ , we have $\sigma(e, v) = f$ and $\sigma(e', v) = f'$. Since $v \sim_i v'$, f and f' belong to some $C_{h'}$, implying (iii). For (iv), one has only to consider a loop γ in A' that intersects more than two of $D_i \times I_{\pm 1}$'s.

($\Leftarrow$) Choose any D_i and $v, v' \in A \cap D_i$ such that $v \neq v'$. Let γ and γ' be (possibly the same) loops in A intersecting D_i at v and v' , respectively. Let e_+, e_-, e'_+ and e'_- denote the edges of Γ corresponding to the components of $A \setminus (\mathcal{D} \times (-1, 1))$ containing $v \times 1$, $v \times -1$, $v' \times 1$ and $v' \times -1$, respectively. Define $v \sim_i v'$ if e_+ and e'_+ happen to be consecutive edges of some C_h ; by (iii), this occurs if and only if $e_- = \sigma(e_+, a_i)$ and $e'_- = \sigma(e'_+, a_i)$ are consecutive in some $C_{h'}$. By (ii), $\sim_i$ is a pairing on $A \cap D_i$. Let A' be obtained from A and from the pair $(\mathcal{D}, \{\sim_i\})$, by the process described in Setting 13. Whenever $v \sim_i v'$ for some i , there is an arc in A' joining $v \times 1$ and $v' \times 1$, and another arc in A' joining $v \times -1$ and $v' \times -1$. Φ sends each loop in A' to some C_h by the definition of $\{\sim_i\}$. Each loop in A' does not intersect $D_i \times I_\epsilon$ more than once for any i and $\epsilon = \pm 1$, by the condition (i). Condition (iv) implies that at least one loop in A' intersects more than two of $D_i \times I_{\pm 1}$'s. Hence, $(\mathcal{D}, \{\sim_i\})$ determines a simple surgery on A . $\square$

For a planar graph $\Gamma \subseteq S^2$, a *bigon neighborhood* of Γ is $\cup_{e \in E(\Gamma)} N_e \subseteq S^2$, where

- (i) each N_e is a 2-cell such that e is properly embedded in N_e , and
- (ii) $N_e \cap N_{e'} = e \cap e'$ for $e \neq e'$.

For example, see Fig. 4.

Definition 18. Consider a 1-submanifold $A = \gamma_1 \cup \dots \cup \gamma_r \subseteq H$ where each γ_i is a loop. If there exists a non-zero integral polynomial of the form $f(x_1, \dots, x_r) = \sum_{i,j \geq 1} c_{ij} x_i^j$ where $c_{ij} \geq 0$ such that $f(A) = \prod_{i,j \geq 1} c_{ij} \gamma_i^j$ admits a simple surgery with respect to a disk structure $\mathcal{D}$ on H , then we say that A *weakly admits a simple surgery (with respect to $\mathcal{D}$)*.

Proposition 19 and 21 are the key steps for the proof of Theorem 22 and 25.




FIGURE 4. An example of a bigon neighborhood. Note that $R'_h \subseteq R_h$.

Proposition 19. *Suppose A is a geometric and diskbusting 1-submanifold of H which intersects a disk structure $\mathcal{D}$ transversely and minimally, such that $W(A, \mathcal{D})$ is minimal. Then A weakly admits a simple surgery with respect to $\mathcal{D}$.*

Proof. By Theorem 12, we may assume that $A \subseteq \partial H$. Write $\mathcal{D} = \{D_1, \dots, D_n\}$ and $\Gamma = W(A, \mathcal{D})$. Let N be a closed regular neighborhood of A in ∂H . $B = \partial N$ denotes the boundary of N as a 2-submanifold of ∂H . Note that $B = A + A$; that is, B consists of two 1-submanifolds each of which is freely homotopic to A . In particular, B intersects $\mathcal{D}$ transversely and minimally. Since A is diskbusting, Γ is connected (Theorem 11); in particular, Γ does not have any isolated vertices. Hence, one can write $S^2 \setminus \Gamma = \coprod_h R_h$ where each R_h is an open disk. Choose a bigon neighborhood $N_1 \subseteq S^2$ of $\Gamma \subseteq S^2$, so that $S^2 \setminus N_1 = \coprod_h R'_h$ for some $R'_h \subseteq R_h$ (Fig. 4). Write $\Gamma' = W(B, \mathcal{D})$. Then we have an edge decomposition $\Gamma' = \cup_h \partial \overline{R'_h} \subseteq S^2$. Define a pairing $\sim_i$ on $B \cap D_i = B \cap \partial D_i$ by $v \sim_i v'$ if v and v' are the endpoints of some interval (denoted as $\alpha_{vv'}$) in $\partial D_i \setminus \text{int}(N)$.

We claim that $(\mathcal{D}, \{\sim_i\})$ determines a simple surgery on B . As in the proof of Proposition 17, choose a product structure on $\mathcal{D} \times [-1, 1]$ such that $B \cap (D_i \times [-1, 1])$ is identified with $(B \cap D_i) \times [-1, 1]$ for each i . In particular, there exist homeomorphisms $D_i \times 0 \rightarrow D_i \times \pm 1$ which are compatible with the product structure. Whenever $v \sim_i v'$, choose $\alpha_{vv'}^+ \subseteq D_i \times 1$ and $\alpha_{vv'}^- \subseteq D_i \times -1$ to be the image of $\alpha_{vv'} \subseteq D_i \times 0$. Let $B' = (B \setminus (\mathcal{D} \times (-1, 1))) \cup (\cup_{v \sim_i v'} \alpha_{vv'}^\pm)$. Then Γ' is obtained from B' by collapsing $\alpha_{vv'}^\pm$ onto the vertices $a_i^{\pm 1}$ for each v and v' in D_i such that $v \sim_i v'$. By this process, each loop in B' collapses onto the cycle $C_h = \partial \overline{R'_h}$ for some h . In the below, let us verify the conditions (i) through (iv) of Proposition 17 with regard to the decomposition $\Gamma' = \cup_h C_h$.

The condition (ii) of Proposition 17 is obvious, since $\partial \overline{R'_h}$ and $\partial \overline{R'_{h'}}$ intersect only at vertices of Γ' . Condition (iii) easily follows from the proof of $(\Rightarrow)$ in Proposition 17. By Theorem 11, Γ is a connected graph without a cut vertex. By Lemma 20, each $\partial \overline{R_h}$ is a simple cycle. This implies $C_h = \partial \overline{R'_h} \approx \partial \overline{R_h}$ is a simple cycle, and hence, we have (i).

The condition (iv) fails only when each $\partial \overline{R'_h}$, and hence each $\partial \overline{R_h}$ also, is a bigon. Since each edge is shared by two regions, the number of edges in Γ must then be the same as the number of regions in $S^2 \setminus \Gamma$. Considering S^2 as a CW-complex having the connected graph Γ as its 1-skeleton, we would have $2 = \chi(S^2) = |V(\Gamma)|$. This contradicts to the fact that the genus of H is larger than 1. $\square$

Lemma 20. *Let $\Gamma \subseteq S^2$ be a connected graph without a cut vertex. If R is a component of $S^2 \setminus \Gamma$, then $\partial \overline{R}$ is a simple closed curve.*

Proof. Let $N(\Gamma)$ denote a closed regular neighborhood of Γ . Denote by T the component of $S^2 \setminus N(\Gamma)$ such that $T \subseteq R$. Since Γ is connected, so is $N(\Gamma)$. Hence, $N(\Gamma)$ is a punctured sphere, and T is an open disk. By the deformation retract $N(\Gamma) \rightarrow \Gamma$, $\partial \overline{T}$ maps to $\partial \overline{R}$. So, $\partial \overline{R}$ is a closed curve and R is an open disk. There exists a polygonal disk $\overline{Q}$ and a quotient map $q: \overline{Q} \rightarrow \overline{R}$ such that $\text{int}(q): Q \rightarrow R$ is a homeomorphism and $\partial q: \partial \overline{Q} \rightarrow \partial \overline{R}$ is a graph map, where $\text{int}(q)$ and ∂q are restrictions of q . We have only to show that ∂q is 1-1. Suppose $q(x) = q(y)$ for some $x \neq y \in (\partial \overline{Q})^{(0)}$. Pick a properly embedded arc $\alpha \subseteq \overline{Q}$ joining x and y , and write $Q \setminus \alpha = Q_1 \cup Q_2$ (Fig. 5 (a)). Since $q(\alpha) \approx S^1$, we can write $S^2 \setminus q(\alpha) = A_1 \cup A_2$ such that A_i is an open disk and $q(Q_i) = R \cap A_i$. So, $q(\alpha)$ separates $q(\partial \overline{Q}_1 \setminus \alpha) \setminus q(x)$ from $q(\partial \overline{Q}_2 \setminus \alpha) \setminus q(y)$ (Fig. 5 (b)). Since $q(\alpha) \cap \Gamma = q(\alpha) \cap \partial \overline{R} = q(x) = q(y)$, $q(x)$ separates Γ . This is a contradiction. $\square$




FIGURE 5. Proof of Lemma 20.

Proposition 21. *Let $U \subseteq F$ be an independent, diskbusting set of root-free, cyclically reduced words. If a 1-submanifold $A \subseteq H$ realizes U with respect to a disk structure $\mathcal{D}$, then the following are equivalent.*

- (1) U is polygonal.
- (2) A weakly admits a simple surgery with respect to $\mathcal{D}$.

Proof. Write $U = \{w_1, \dots, w_r\}$ and $A = \gamma_1 \cup \dots \cup \gamma_r$ so that γ_i realizes w_i with respect to $\mathcal{D}$. Recall our convention that A intersects $\mathcal{D}$ transversely and minimally (Remark 10).

(1) $\Rightarrow$ (2): Suppose there exists a U -polygonal surface S and an associated immersion $\phi: S^{(1)} \rightarrow \text{Cay}(F)/F$ satisfying $\chi(S) < m(S)$. For $i, j \geq 1$, let c_{ij} denote the number of polygonal disks P on S such that the composition of immersions $\partial P \rightarrow S^{(1)} \xrightarrow{\phi} \text{Cay}(F)/F$ reads w_i^j . In particular, $m(S) = \sum_{i,j} c_{ij}$. One can write $S = \coprod_i \coprod_j \coprod_{k=1}^{c_{ij}} P_{ijk} / \sim$ for some side-pairing $\sim$ such that each P_{ijk} is a polygonal disk whose boundary reads w_i^j . Put $f(x_1, \dots, x_r) = \sum_{i,j} c_{ij} x_i^j$ and $\Gamma = W(f(U))$. Let σ denote the connecting map associated with Γ . By Remark 9 (2), there is a natural 1-1 correspondence ρ between the edges of Γ and the corners (i.e., two

adjacent edges) of $\coprod_{i,j,k} P_{ijk}$. For instance, a vertex $v \in \partial P_{ijk}$ at which an a_1 -edge is incoming and an a_2 -edge is outgoing corresponds to an edge of Γ joining a_1 and a_2^{-1} (Fig. 6).




FIGURE 6. The 1 – 1 correspondence ρ .

Write $S^{(0)} = \{v_1, \dots, v_t\}$. $\text{LINK}(v_h) \approx S^1$ denotes the link of v_h in S . ρ^{-1} maps each $\text{LINK}(v_h)$ to some cycle $C_h \subseteq \Gamma$. Since $\phi: S^{(1)} \rightarrow \text{Cay}(F)/F$ is locally injective, each C_h is a simple cycle. This proves the condition (i) of Proposition 17. The condition (ii) follows from the fact that ρ is a 1 – 1 correspondence.

Suppose e and e' are consecutive edges in some C_h . e and e' correspond to an adjacent pair of edges (corners) in $\text{LINK}(v_h)$. Without loss of generality, assume $a_q \in \partial e \cap \partial e' \subseteq V(\Gamma)$, and let $v_{h'} \in S^{(0)}$ be the other endpoint of the a_q -edge incoming at v_h (Fig. 7 (a)). $\rho(e)$ is the corner of some P_{ijk} at v_h , and similarly, $\rho(e')$ is the corner of some $P_{i'j'k'}$ at v_h . From the definition of σ , one can see that the corners of P_{ijk} and $P_{i'j'k'}$ at $v_{h'}$ correspond to $f = \sigma(e, a_q)$ and $f' = \sigma(e', a_q)$, respectively. f and f' belong to some $C_{h'}$ as consecutive edges (Fig. 7 (b)). This proves the condition (iii) of Proposition 17.




FIGURE 7. Visualizing the connecting map σ .

Now we will use the assumption $\chi(S) < m(S)$ as follows. The number of vertices, edges and faces in $S = \coprod_{i,j,k} P_{ijk} / \sim$ are t , $\sum_{i,j} c_{ij} |w_i^j|/2$ and $\sum_{i,j} c_{ij}$, respectively. Hence,

$$\begin{aligned} \sum_{i,j} c_{ij} = m(S) &> \chi(S) = t - \sum_{i,j} c_{ij} |w_i^j|/2 + \sum_{i,j} c_{ij}, \\ 2t &< \sum_{i,j} c_{ij} |w_i^j| = |E(\Gamma)| = \sum_{h=1}^t |E(C_h)|. \end{aligned}$$

It follows that $|E(C_h)| > 2$ for some h ; that is, the condition (iv) of Proposition 17 is satisfied.

(2) $\Rightarrow$ (1): Suppose $f(x_1, \dots, x_r) = \sum_{i,j \geq 1} c_{ij} x_i^j$ is a non-zero integral polynomial with $c_{ij} \geq 0$ such that there is a simple surgery on $f(A) = f(\gamma_1, \dots, \gamma_r)$. Write $\Gamma = W(f(A), \mathcal{D}) = \cup_{h=1}^t C_h$ so that the conditions in Proposition 17 are satisfied. For each i, j and $1 \leq k \leq c_{ij}$, take a polygonal disk P_{ijk} equipped with an immersion $\partial P_{ijk} \rightarrow \text{Cay}(F)/F$ reading w_i^j ; so, each edge in $\coprod_{i,j,k} \partial P_{ijk}$ carries a label by $\mathcal{S} = \{a_1, \dots, a_n\}$ as well as an orientation induced from the orientations of the edges in $\text{Cay}(F)/F$. There is a 1 – 1 correspondence ρ between the edges of Γ and the corners of $\coprod_{i,j,k} P_{ijk}$ as in the proof of (1) $\Rightarrow$ (2). Let $1 \leq q \leq n$, and consider two a_q -edges $x \in E(\partial P_{ijk})$ and $x' \in \partial E(P_{i'j'k'})$. We declare that $x \sim x'$ if ρ^{-1} sends the corners at the terminal vertices of x and x' to consecutive edges e and e' in some C_h (Fig. 7 (a)). By the condition (ii) of Proposition 17, $e \in E(\Gamma)$ and $a_q \in \partial e$ uniquely determines $e' \in E(\Gamma)$ such that e and e' are consecutive at a_q in some C_h . By the condition (iii), the other corners of x and x' correspond to consecutive edges $f = \sigma(e, a_q)$ and $f' = \sigma(e', a_q)$ in some $C_{h'}$. Hence, $\sim$ defines a side-pairing on $\coprod_{i,j,k} P_{ijk}$ that matches the labels and the orientations of the edges; moreover, ρ determines a 1 – 1 correspondence between the links of the vertices in the closed surface $S = \coprod_{i,j,k} P_{ijk} / \sim$ and the cycles $C_1, C_2, \dots, C_t$. Since each C_h is simple, there is an immersion $S^{(1)} \rightarrow \text{Cay}(F)/F$ induced by the given immersion $\coprod_{i,j,k} \partial P_{ijk} \rightarrow \text{Cay}(F)/F$. Note that $m(S) = \sum_{i,j} c_{ij}$. As in the proof of (1) $\Rightarrow$ (2),

$$\chi(S) = t - |E(\Gamma)|/2 + m(S) = t - \sum_{h=1}^t |E(C_h)|/2 + m(S).$$

By the condition (iv) of Proposition 17, $|E(C_h)| > 2$ for some h . Therefore, $\chi(S) < m(S)$. $\square$

Theorem 22. *Suppose $U \subseteq F$ is an independent, diskbusting set of root-free, cyclically reduced words. If U is geometric, then U is equivalent to a polygonal set of words in F .*

Proof. We may assume that U is minimal, by applying an automorphism of F if necessary. Let $U = \{w_1, \dots, w_r\}$ be realized by $A \subseteq H$ with respect to a disk structure $\mathcal{D}$. By Theorem 12, we can choose A so that $A \subseteq \partial H$ and $W(A, \mathcal{D})$ is minimal. Proposition 19 implies that A weakly admits a simple surgery with respect to $\mathcal{D}$. Hence, U is polygonal by Proposition 21. $\square$

Now we consider virtually polygonal words (Definition 3). Note that polygonal words are virtually polygonal. The converse is not true. Actually, there exists a non-polygonal word which becomes polygonal only after an automorphism of F is applied. An example given in [7] is $w = abab^2ab^3 \in F_2 = \langle a, b \rangle$. While an elementary argument shows that w is not polygonal, the automorphism ϕ defined by $\phi(a) = ab^{-2}$ and $\phi(b) = b$ maps w to a Baumslag–Solitar relator $w' = a(a^2)^b$. Any Baumslag–Solitar relator $a^p(a^q)^b$ is polygonal for $pq \neq 0$ [7].

Definition 23. A 1-submanifold $A \subseteq H$ *virtually admits a simple surgery* if there exists a finite cover $p : H' \rightarrow H$ such that $p^{-1}(A) \subseteq H'$ weakly admits a simple surgery.

Proposition 24. *Let $w \in F$ be a root-free and diskbusting word realized by a loop $\gamma \subseteq H$. Then w is virtually polygonal if and only if γ virtually admits a simple surgery.*

Proof. Let F' be a finite-index subgroup of F and $\mathcal{S}'$ be a free basis for F' . Denote by $\hat{w}_{F'}$ a transversal for $\underline{w}_{F'}$, so that $\hat{w}_{F'}$ is an independent set of cyclically reduced words written in $\mathcal{S}'$. $\hat{w}_{F'}$ is realized by $p^{-1}(\gamma)$ with respect to some disk structure $\mathcal{D}'$ on H' . By Proposition 21, $\hat{w}_{F'}$ is polygonal with respect to $\mathcal{S}'$ if and only if $p^{-1}(\gamma)$ weakly admits a simple surgery with respect to $\mathcal{D}'$. $\square$

Theorem 25. *A diskbusting and virtually geometric word in F is virtually polygonal.*

Proof. Let w be diskbusting and virtually geometric. We may assume w is root-free and cyclically reduced. Let $\gamma \subseteq H$ realize w with respect to a given disk structure $\mathcal{D}$. Suppose $p : H' \rightarrow H$ is a finite cover such that $p^{-1}(\gamma)$ is freely homotopic to $A \subseteq \partial H'$. It is elementary to see that A is also diskbusting in H' . By Proposition 19, A weakly admits a simple surgery with respect to some disk structure $\mathcal{D}'$ on H' . By Proposition 24, w is virtually polygonal. $\square$

Theorem 26 underlines the importance of virtual polygonality.

Theorem 26. *If a root-free word $w \in F$ is virtually polygonal, then $D(w)$ contains a hyperbolic surface group.*

Proof. We may assume w is cyclically reduced by applying an automorphism of $F = \langle \mathcal{S} \rangle$ if necessary. Let $\gamma_w \subseteq \text{Cay}(F)/F$ read w . Let F' be a finite-index subgroup of F such that a transversal $\hat{w}_{F'}$ for $\underline{w}_{F'}$ is polygonal with respect to some free basis $\mathcal{S}'$ for F' . Recall that $X_{\mathcal{S}'}(\hat{w}_{F'})$ denote the 2-dimensional CW-complex obtained by taking two copies of $\text{Cay}_{\mathcal{S}'}(F')/F'$ and gluing cylinders along the loops reading $\hat{w}_{F'}$ considered as a set of words written in $\mathcal{S}'$. By Theorem 6, $D(\hat{w}_{F'}) = \pi_1(X_{\mathcal{S}'}(\hat{w}_{F'}))$ contains a hyperbolic surface group. As was introduced in Section 2, $Y(w, F')$ denotes the finite cover of $X_{\mathcal{S}}(w)$ obtained by taking two copies of $\text{Cay}(F)/F'$ and gluing cylinders along the copies of the elevations of γ_w . Since the homotopy equivalence $\text{Cay}(F)/F' \rightarrow \text{Cay}_{\mathcal{S}'}(F')/F'$ maps each elevation of γ_w to a loop realizing an element in $\hat{w}_{F'}$, we have a homotopy equivalence $Y(w, F') \simeq X_{\mathcal{S}'}(\hat{w}_{F'})$. Hence $D(w) \geq \pi_1(Y(w, F')) = D(\hat{w}_{F'})$ and so, $D(w)$ contains a hyperbolic surface group. $\square$

Corollary 27 (Gordon–Wilton [6]). *If $w \in F$ is root-free, virtually geometric and diskbusting, then $D(w)$ contains a hyperbolic surface group.* $\square$

Converses of Theorem 22 and 25 do not hold. Actually, we will prove the following theorem in Section 4.

Theorem 28. *There exist polygonal words which are not virtually geometric.*

Virtual geometricity does not imply geometricity, in general. For example, $w = a^2b^{-1}ab \in F_2 = \langle a, b \rangle$ is virtually geometric, but not geometric [6]. It is not known whether virtual polygonality is strictly weaker than polygonality up to $\text{Aut}(F)$.

Question 29. Is a virtually polygonal word equivalent to a polygonal word?

Tiling Conjecture ([7]). *A minimal diskbusting word in F is polygonal.*


While the Tiling Conjecture is not resolved yet, we propose a weaker conjecture.

Virtual Tiling Conjecture. *A diskbusting word in F is virtually polygonal.*

Remark 30. Let $w \in F$ be a diskbusting and root-free word realized by $\gamma \subseteq H$. Consider the following hypotheses on w and γ :

- $*$: No further hypothesis on w .
- (V)Geom : w is (virtually) geometric.
- WSS : γ weakly admits a simple surgery.
- VSS : γ virtually admits a simple surgery.
- EPoly: w is equivalent to a polygonal word.
- VPoly : w is virtually polygonal.
- DSurf : $D(w)$ contains a hyperbolic surface group.

Note that none of the above hypotheses are dependent on the choice of a free basis for F , or equivalently, on the choice of a disk structure on H . We can summarize the content of this section as the following diagram:



where each dashed arrow is a question or conjecture unanswered in this paper, and each broken arrow with $\times$ in the middle means a false implication. Question 29, the Tiling Conjecture and the Virtual Tiling Conjecture are equivalent to (A), (B) and (C), respectively. (D) is equivalent to the Gromov's conjecture for the groups of the form $D(w)$.

Let us set $F_3 = \langle a, b, c \rangle$ and $F_4 = \langle a, b, c, d \rangle$. Throughout this section, put $w_1 = bbaaccabc \in F_3$ and $w_2 = aabbacbccadbdcd \in F_4$. A graph Γ is k -valent if the valence of each vertex is k . Γ is k -edge-connected if one cannot disconnect Γ by removing the interior of $k - 1$ or fewer edges. In [8], Manning proved (a stronger version of) the following theorem.

Theorem 31 ([8]). *Let $w \in F$ be a cyclically reduced word such that for some $k \geq 3$, $W(w)$ is a k -valent, k -edge-connected and non-planar graph. Then w is not virtually geometric.* $\square$

An example of graphs satisfying the hypothesis in Theorem 31 is the complete bipartite graph $K_{k,k}$ for $k \geq 3$. Manning noted that $W(w_1)$ is $K_{3,3}$, and hence, w_1 is not virtually geometric [8]. Since $W(w_2)$ is $K_{4,4}$, w_2 is not virtually geometric, either. On the other hand,

Proposition 32. (1) $w_1 = bbaaccabc$ is polygonal.
(2) $w_2 = aabbacbccadbcd$ is polygonal.

Proof. (1) Let P_1 be a polygonal disk whose boundary reads w_1^2 . Name the edges of ∂P_1 as $1, 2, \dots, 18$ so that the edge named by i corresponds to the i -th letter of w_1^2 . Define a side-pairing $\sim_1$ on P_1 as shown in Fig. 8 (a). That is to say, the edges of ∂P_1 are paired by $\sim_1$ as

$$\{1, 2\}, \{3, 7\}, \{4, 16\}, \{5, 15\}, \{6, 18\}, \{8, 11\}, \{9, 14\}, \{10, 17\}, \{12, 13\}.$$

A neighborhood of each vertex in $S_1 = P_1/\sim_1$ is illustrated in Fig. 8 (b). One sees that no two incoming edges (or two outgoing edges) of the same label exist at each vertex. Therefore, S_1 is a w_1 -polygonal surface. Since $\chi(S_1) = |S^{(0)}| - |S^{(1)}| + |S^{(2)}| = 4 - 18/2 + 1 = -4 < 1$, we conclude that w_1 is polygonal.

(2) The proof that w_2 is polygonal is almost identical with (1) by considering a side-pairing $\sim_2$ on P_2 , where P_2 is a polygonal disk whose boundary reads w_2 ; see Fig. 9 (a). Specifically, $\sim_2$ identifies the edges of ∂P_2 as

$$\{1, 2\}, \{3, 12\}, \{4, 7\}, \{5, 10\}, \{6, 14\}, \{8, 9\}, \{11, 16\}, \{13, 15\}.$$

One again sees that $S_2 = P_2/\sim_2$ is a w_2 -polygonal surface, from the description of the links in Fig. 9 (b). Since $\chi(S_2) = 4 - 16/2 + 1 = -3 < 1$, w_2 is polygonal. $\square$

Remark 33. (1) Let Γ be a graph immersed in $\text{Cay}(F)/F$. Then Γ embeds into $\text{Cay}(F)/F'$ for some $[F : F'] < \infty$ such that $|\text{Cay}(F)/F'^{(0)}| = |\Gamma^{(0)}|$. In particular, the degree of the cover $\text{Cay}(F)/F' \rightarrow \text{Cay}(F)/F$ can be chosen to be $|\Gamma^{(0)}|$; see [11]. By the proof of Theorem 6, we observe that if $w \in F$ has a closed w -polygonal surface S , then $X(w)$ has a finite cover of degree $|S^{(0)}|$ containing a closed surface S' such that $\chi(S') = 2(\chi(S) - m(S))$. From the proof of Proposition 32 and Fig. 8, we see that $X(w_1)$ has a finite cover of degree 4 that contains a closed surface of Euler characteristic $2(\chi(S_1) - 1) = -10$. Similarly, a degree-4 cover of $X(w_2)$ contains a closed surface of Euler characteristic $2(\chi(S_2) - 1) = -8$.

(2) Let $\Gamma_1 = W(w_1^2)$ and $\Gamma_2 = W(w_2)$. S_1 has one vertex of valence six and three vertices of valence four. By the proof of Proposition 17 and 21, Γ_1 can be written as the union of one simple cycle of length six and three simple cycles of length four. Also, Γ_2 is the union of four simple cycles of length four.





FIGURE 8. Proposition 32 (1). Single, double and triple arrows denote a -, b - and c -edges, respectively.

Acknowledgement. The author appreciates Cameron Gordon for inspirational conversations. The author is grateful to Henry Wilton for helpful discussion on Section 2.1, and to Alan Reid for his guidance through this work.

(a) A side-pairing $\sim_2$ on P_2 .(b) Vertices in $S_2 = P_2/\sim_2$.FIGURE 9. Proposition 32 (2). Single, double, triple, and white arrows denote a -, b -, c - and d -edges, respectively.

REFERENCES

- [1] J Berge. Heegaard documentation. Preprint (c.1990), <http://www.math.uic.edu/~t3m>.
- [2] M Bestvina. Questions in geometric group theory. <http://www.math.utah.edu/bestvina>.
- [3] M Bestvina and M Feighn. A combination theorem for negatively curved groups. *J. Differential Geom.*, 35(1):85–101, 1992.
- [4] D Calegari. Surface subgroups from homology. *Geometry & Topology*, 12(4):1995–2007, 2008.
- [5] R D Canary. Ends of hyperbolic 3-manifolds. *J. Amer. Math. Soc.*, 6(1):1–35, 1993.
- [6] C McA Gordon and H Wilton. On surface subgroups of doubles of free groups. Preprint, arXiv:0902.3693v1.
- [7] S Kim and H Wilton. On polygonality of words in free groups. Preprint, arXiv:0910.4709v1.
- [8] J F Manning. Virtually geometric words and Whitehead’s algorithm. Preprint, arXiv:0904.0724v2.
- [9] P Scott. Subgroups of surface groups are almost geometric. *J. London Math. Soc. (2)*, 17(3):555–565, 1978.
- [10] J R Stallings. Whitehead graphs on handlebodies. *Geometric group theory down under (Canberra, 1996)*, 317–330, de Gruyter, Berlin, 1999.
- [11] J R Stallings. Topology of finite graphs. *Invent. Math.*, 71(3):551–565, 1983.
- [12] R Stong. Diskbusting elements of the free group. *Math. Res. Lett.*, 4(2-3):201–210, 1997.
- [13] J H C Whitehead. On certain sets of elements in a free group. *Proceedings of the London Mathematical Society*, s 2-41(1):48, Jan 1936.
- [14] Henry Wilton. Hall’s theorem for limit groups. *Geom. Funct. Anal.*, 18(1):271–303, 2008.
- [15] D T Wise. Subgroup separability of graphs of free groups with cyclic edge groups. *Q. J. Math.*, 51(1):107–129, 2000.
- [16] H Zieschang. Simple path systems on full pretzels. *Mat. Sb. (N.S.)*, 66 (108):230–239, 1965. Translated in *AMS Translations (2)*, 92: 127–137, 1970.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN
 E-mail address: shkim@math.utexas.edu